

References (76)

1. Schuller D, Cardoso F, Sousa S, Gomes P, Gomes AC, Santos MAS, Casal M. Genetic diversity and population structure of *Saccharomyces cerevisiae* strains isolated from different grape varieties and winemaking regions PLOSone. 2012; 7(2):e32507
2. Tello J, Cordero-Bueso G, Aporta I, Cabellos JM, Arroyo T. Genetic diversity in commercial wineries: effect of the farming system and vinification management on wine yeasts. J Appl Microbiol. 2011; 112:302-15
3. Boulton RB, Singleton VL, Bisson LF, Kunkee RE. Principles and Practices of Winemaking. New York: Chapman & Hall; 1996
4. Bokulich N, Ohta M, Richardson PM, Mills DA. Monitoring seasonal changes in winery resident microbiota. PLOSone 2013; 8(6) e66437
5. Bokulich NA, Thorngate JH, Richardson PM, Mills DA. Microbial biogeography of wine grapes is conditioned by cultivar, vintage and climate. Proc Natl Acad Sci USA. 2014; 111:e139-e148
6. Sacchi K L, Bisson L F, Adams DO. A review of the effect of winemaking techniques on phenolic extraction in red wines. Am J Enol Vitic. 2005; 56:197-206
7. Fleet GH. Wine yeasts for the future. FEMS Yeast Res. 2008; 8:979-95
8. Hierro N, Gonzalez A, Mas A, Guillamon JM. Diversity and evolution of non-*Saccharomyces* yeast populations during wine fermentation: effect of grape ripeness and cold maceration. FEMS Yeast Res. 2006; 6:102-11
9. Zott K, Miot-Sertier C, Claisse O, Lonvaud-Funel A, Masneuf-Pomarede I. Dynamics and diversity of non-*Saccharomyces* yeasts during the early stages in winemaking. Int J Food Microbiol. 2008; 125:197-203
10. Osborne JP, Edwards CG. Bacteria in winemaking. Adv Food Nutr Res. 2005; 50:139-77
11. Bartowsky E, Henschke PA. Acetic acid bacteria spoilage of bottled wine – a review. Int J Food Microbiol. 2008; 125:60-70.
12. Belloch C, Perez-Torrado R, Gonzalez SS, Perez-Ortin JE, Garcia-Martinez J, Querol A, Barrio E. Chimeric genomes of natural hybrids of *Saccharomyces cerevisiae* and *Saccharomyces kudriavzevii*, Appl Environ Microbiol. 2009; 75: 2534-2544
13. Bisson LF. Geographic origin and diversity of wine strains of Am J Enol Vitic. 2012; 63:165-176
14. Dequin S, Casaregola S. The genomes of fermentative *Saccharomyces*. C R Biologies. 2011; doi: 10.1016/j.crvi.2011.05.019
15. González SS, Barrio E, Gafner J, Querol A. Natural hybrids from *Saccharomyces cerevisiae*, *Saccharomyces bayanus* and *Saccharomyces kudriavzevii* in wine fermentations. FEMS Yeast Res. 2006; 6:1221-34
16. Gonzalez SS, Barrio E, Querol A. Molecular characterization of new natural hybrids of *Saccharomyces cerevisiae* and *kudriavzevii* in brewing. Appl Envion Microbiol. 2008; 74:2314-20
17. Masneuf I, Hansen J, Groth C, Piskur J, Dubourdieu D. New hybrids between *Saccharomyces sensu stricto* yeast species found among wine and cider production strains. Appl Environ Microbiol. 1998; 64:3887-92
18. Naumova, E. S., Naumov, G. I., Masneuf-Pomarede, I., Aigle, M. and Dubourdieu, D. Molecular genetic study of introgression between *Saccharomyces bayanus* and *S. cerevisiae*. Yeast. 2005; 22:1099-115
19. Bisson LF. Stuck and sluggish fermentations. Am J Enol Vitic. 1999; 50:107-119
20. Baumes R. Wine aroma precursors. In: Moreno-Arribas MV, Polo MC, editors. Wine Chemistry and Biochemistry. New York. Springer Science + Business Media; 2009
21. Ferreira V, Lopez R, Escudero A, Cacho JF. The aroma of Grenache red wine: hierarchy and nature of its main odorants. J Sci Food Agric. 1998; 77:259-67
22. Ferreira V, Cacho J. Identification of impact odorants of wines. In: Moreno-Arribas MV, Polo MC, editors. Wine Chemistry and Biochemistry. New York. Springer Science + Business Media; 2009
23. Guth H. Identification of character impact odorants of different white wine varieties. J Agric Food Chem. 1997; 45:3022-26
24. Guth H. Quantitation and sensory studies of character impact odorants of different white wine varieties. J Agric Food Chem. 1997; 45:3027-32
25. Lee SH, Seo M, Riu M, Cotta JP, Block DE, et al. Vine microclimate and norisoprenoid concentration in Cabernet Sauvignon grapes and wines. Am J Enol Vitic. 2007; 58:291-301
26. Mendes-Pinto MM. Carotenoid breakdown products the – norisoprenoids – in wine aroma. Arch Biochem Biophys. 2009; 483:236-45

27. Heard GM, Fleet GH. Growth of natural yeast flora during the fermentation of inoculated wines. *Appl Environ Microbiol.* 1985; 50:727-28
28. Polaskova P, Herszag J, Ebeler E. Wine flavour: chemistry in a glass. *Chem Soc Rev.* 2008; 37:2478- 89
29. Preston LD, Block DE, Heymann H, Soleas G, Noble AC, et al. Defining vegetal aromas in Cabernet Sauvignon using sensory and chemical evaluations. *Am J Enol Vitic.* 2008; 59:137-45
30. Rapp A, Mandery H. Wine aroma. *Experientia* 1986; 42:873-84
31. Tominaga T, Peyrot des Gachons C, Dubourdieu D. A new type of flavour precursors in *Vitis vinifera* L. cv. Sauvignon Blanc: S-cysteine conjugates. *J Agric Food Chem.* 1998; 46:5215-19
32. Tominaga T, Baltenweck Guyot R, Peyrot des Gachons C, Dubourdieu D. Contribution of volatile thiols to the aromas of white wines made from several *Vitis vinifera* grape varieties. *Am J Enol Vitic.* 2000; 51:178-81
33. Tominaga T, Blanchard L, Darriet P and Dubourdieu D. A powerful aromatic volatile thiol, 2-furanmethanethiol, exhibiting roast coffee aroma in wines made from several *Vitis vinifera* grape varieties. *J Agric Food Chem.* 2009; 48:1799-802
34. Vilanova M, Cortes S, Santiago JL, Martinez C, Fernandez E. Aromatic compounds in wines produced during fermentation: effect of three red cultivars. *Int J Food Prop.* 2007; 10:867-75
35. Zhang M, Xu Q, Duan C, Qu W, Wu Y. Comparative study of aromatic compounds in young red wines from Cabernet Sauvignon, Cabernet Franc and Cabernet Gernischet varieties in China. *J Food Sci.* 2007; 72:248-52
36. Pozo-Bayon MA, Reineccius G. Interactions between wine matrix macro-components and aroma compounds. In: Moreno-Arribas MV, Polo MC, editors. *Wine Chemistry and Biochemistry.* New York. Springer Science + Business Media; 2009
37. Ugliano M, Henschke P. Yeasts and Wine Flavour. In: Moreno-Arribas MV, Polo MC. Editors. *Wine Chemistry and Biochemistry.* New York. Springer Science + Business Media; 2009
38. Cullere L, Cacho J, Ferreira V. An assessment of the role played by some oxidation-related aldehydes in wine aroma. *J Agric Food Chem.* 2007; 55:876-81
39. Cutzach I, Chatonnet P, Dubourdieu D. Influence of storage conditions on the formation of some volatile compounds in white fortified wines (*Vins doux naturels*) during the aging process. *J Agric Food Chem.* 2009; 48:2340-45
40. Jarauta I, Cacho J, Ferreira V. Concurrent phenomena contributing to the formation of the aroma of wine during aging in oak wood: an analytical study. *J Agric Food Chem.* 2005; 53:4166-77
41. Spillman PJ, Pollnitz AP, Liacopoulos D, Pardon KH, Sefton MA. Formation and degradation of furfuryl alcohol, 5-methylfurfuryl alcohol, vanillyl alcohol and their ethyl esters in barrel-aged wines. *J Agric Food Chem.* 1998; 46:657-63
42. Bardi L, Crivelli C, Marzona M. Esterase activity and release of ethyl esters of medium-chain fatty acids by *Saccharomyces cerevisiae* during anaerobic growth. *Can J Microbiol.* 1998; 44:1171-76
43. Bartowsky EJ, Pretorius IS. Microbial formation and modification of flavour and off-flavour compounds in wine. In: Konig H, Unden G, Frohlich J, editors. *Biology of Microorganisms on Grapes, in Musts and in Wine.* Berlin Heidelberg. Springer-Verlag; 2009
44. Dickinson JR, Eshantha L, Salgado J, Hewlins MJE. The catabolism of amino acids to long chain and complex alcohols in *Saccharomyces cerevisiae*. *J Biol Chem.* 2003; 278:8028-34
45. Dubourdieu D, Tominaga T, Masneuf I, Peyrot des Gachons C, Murat ML. The role of yeasts in grape flavour development during fermentation: the example of Sauvignon blanc. *Am J Enol Vitic.* 2006; 57:81-8
46. Fernandez-Gonzalez M, Di Stefano R. Fractionation of glycoside aroma precursors in neutral grapes. Hydrolysis and conversion by *Saccharomyces cerevisiae*. *Lebensm-Wiss. U. Technol.* 2003; 37:467-73
47. Ferreira V, Lopez R, Aznar M. Olfactometry and aroma extract dilution analysis of wines. In: Jackson JF, Linskens HF, editors. *Analysis of Taste and Aroma.* Berlin Heidelberg. Springer-Verlag; 2009
48. Hazelwood LA, Daran JM, van Maris AJA, Pronk JT, Dickinson JR. The Ehrlich pathway for fusel alcohol production: a century of research on *Saccharomyces cerevisiae* *Appl Environ Microbiol.* 2008; 74:2259-66
49. Howell KS, Klein M, Swiegers JH, Hayasaki Y, Elsey GM, et al. Genetic determinants of volatile thiol release by *Saccharomyces cerevisiae* during wine fermentation. *Appl Environ Microbiol.* 2005; 71:5420-26
50. Howell KS, Cozzolino, Bartowsky EJ, Fleet GH, Henschke PA. Metabolic profiling as a tool for revealing *Saccharomyces* interactions during wine fermentation. *FEMS Yeast Res.* 2006; 6:91-101

51. Kotseridis Y, Baumes R. Identification of impact odorants in Bordeaux red grape juice, in the commercial yeast used for its fermentation, and in the produced wine. *J Agric Food Chem.* 2000; 48:400-6
52. Landaud S, Helinck S, Bonnarme P. Formation of volatile sulfur compounds and metabolism of methionine and other sulfur compounds in fermented foods. *Appl Microbiol Biotechnol.* 2008; 77:1191-205
53. Marchand S, de Revel G, Bertrand A. Approaches to wine aroma: Release of aroma compounds from reactions between cysteine and carbonyl compounds in wine. *J Agric Food Chem.* 2000; 48:4890-95
54. Mason AB, Dufour JP. Alcohol acetyltransferases and the significance of ester synthesis in yeast. *Yeast.* 2000; 16:1287-98
55. Miller AC, Wolff SR, Bisson LF, Ebeler SE. Yeast strain and nitrogen supplementation: dynamics of volatile ester formation in Chardonnay juice fermentations. *Am J Enol Vitic.* 2007; 58:470-83
56. Molina AM, Swiegers JH, Varela C, Pretorius IS, Agosin E. Influence of wine fermentation temperature on the synthesis of yeast-derived volatile aroma compounds. *Appl Microbiol Biotechnol.* 2007; 77:675-87
57. Moreira N, Mendes F, Pereira O, Guedes de Pinho P, Hogg T, et al. Volatile sulphur compounds in wines related to yeast metabolism and nitrogen composition of grape musts. *Anal Chim Acta* 2002; 458:157-67
58. Murat ML, Masneuf I, Darriet P, Lavigne V, Tominaga T, et al. Effect of *Saccharomyces cerevisiae* yeast strains on the liberation of volatile thiols in Sauvignon blanc wine. *Am J Enol Vitic.* 2001; 52:136-9
59. Nykanen L. Formation and occurrence of flavour compounds in wine and distilled beverages. *Am J Enol Vitic.* 1986; 37:84-96
60. Parkkinen E, Suomalainen H. Esterases of Baker's yeast. II. Substrate specificities towards esters formed during sugar fermentations. *J Inst Brew.* 1982; 88:34-8
61. Rauhut D. Yeasts – Production of sulfur compounds. In: Fleet GH, editor. *Wine Microbiology and Biotechnology.* Chur. Harwood Academic Publishers GmbH; 1993
62. Rauhut D. Usage and formation of sulphur compounds. In: Konig H, Unden G, Frohlich J, editors. *Biology of Microorganisms on Grapes, in Musts and in Wine.* Berlin Heidelberg. Springer-Verlag; 2009
63. Romano P, Fiore C, Paraggio M, Caruso M, Capece A. Function of yeast species and strains in wine flavour. *Int J Food Microbiol.* 2003; 86:169-80
64. Rossouw D, Naes T, Bauer F. Linking gene regulation and the exo-metabolome: A comparative transcriptomics approach to identify genes that impact the production of volatile aroma compounds in yeast. *BMC Genomics.* 2008; doi:10.1186/1471-2164-9-530
65. Saerens SM, Delvaux FR, Verstrepen KJ, Thevelein JM. Production and biological function of volatile esters in *Saccharomyces cerevisiae*. *Microbial Biotech.* 2009; doi:1111/j.1751-7915.2009.00106
66. Schoondermark-Stolk SA, Jansen M, Verkleij AJ, Verrrips CT, Euverink GJW, et al. Genome-wide transcription survey on flavour production in *Saccharomyces cerevisiae*. *World J Microbiol Biotechnol.* 2006; 22:1347-56
67. Schoondermark-Stolk SA, Jansen M, Veurink JH, Verkleij AJ, Verrrips CT, et al. Rapid identification of target genes for 3-methyl-1-butanol production in *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol.* 2006; 70:237-46.
68. Spiropoulos A, Tanaka J, Flerianos I, Bisson LF. Characterization of hydrogen sulfide formation in commercial and natural wine isolates of *Am J Enol Vitic.* 2000; 51:233-48
69. Suomalainen H. Yeast esterases and aroma esters in alcoholic beverages. *J Inst Brew.* 1981; 87:296-300
70. Swiegers JH, Bartowsky EJ, Henschke PA, Pretorius IS. Yeast and bacteria modulation of wine aroma and flavour. *Aus J Grape Wine Res.* 2005; 11:139-73
71. Swiegers JH, Pretorius IS. Modulation of volatile sulfur compounds by wine yeast. *Appl Microbiol Biotechnol.* 2007; 74:954-60
72. Thorngate JH. Yeast strain and wine flavour: Nature or nurture? In: Waterhouse AL, Ebeler SE, editors. *Chemistry of Wine Flavour.* Washington DC. American Chemical Society; 1998
73. Thurston PA, Quain DE, Tibb RS. Lipid metabolism and the regulation of volatile ester synthesis in *Saccharomyces cerevisiae*. *J Inst Brew* 1982; 88:90-4
74. Ugliano M, Bartowsky EJ, McCarthy J, Moio L, Henschke PA. Hydrolysis and transformation of grape glycosidically bound volatile compounds during fermentation with three *Saccharomyces* yeast strains. *J Agric Food Chem.* 2006; 54:6322-31

75. Verstrepen KJ, Van Laere SDM, Vanderhaegen BMP, Derdelinckx G, Dufour JP, et al. Expression levels of the yeast alcohol acetyltransferase genes ATF1, Lg-ATF1, and ATF2 control the formation of a broad range of volatile esters. *Appl Environ Microbiol*. 2003; 69:5228-37
76. Yoshimoto H, Fukushige T, Yonezawa T, Sone H. Genetical and physiological analysis of branched-chain alcohols and isoamyl acetate production in *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol*. 2002; 59:501-8